Mail Stop Amendment

RECEIVED
CENTRAL FAX CENTER

DEC 2 1 2005

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

DN A01195

In re application of:

Jacobson, et.al.

Serial No.:

10/645,427

: Group Art Unit:

1617

Filed:

08/21/03

: Examiner:

S. Clardy

For:

A Method to Inhibit Ethylene Responses in Plants

Mail Stop Amendment Commissioner for Patents P. O. Box 1450 Alexandria, VA 22313-1450

CERTIFICATION OF FACSIMILE TRANSMISSION

I hereby certify that the following papers are being facsimile transmitted to the Patent and Trademark Office on the date shown below.

RESPONSE

December 21, 2005	Thomas D. Rogenson	
Date	Signature	

Total Pages 3

Fax No. 571-273-8300

Mail Stop Amendment

RECEIVED
CENTRAL FAX CENTER

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

ROHM & HAAS

DEC 2 1 2005

DN A01195

In re application of:

Jacobson, et.al.

Serial No.:

10/645,427

: Group Art Unit:

1617

Filed:

08/21/03

: Examiner:

S. Clardy

For:

A Method to Inhibit Ethylene Responses in Plants

Mail Stop Amendment Commissioner for Patents P. O. Box 1450 Alexandria, VA 22313-1450 Dear Sir:

RESPONSE

This is in response to the Office Action dated September 23, 2005 for the aboveidentified application.

REMARKS

Claims 1-8 are pending. Claims 1-8 are rejected.

Rejection under 35 USC §103(a)

Claims 1-8 are rejected under 35 USC §103(a) as being unpatentable over the combined teachings of Daly et al (US 6,017,849) and Sisler (US 6,65,549). Daly teaches substituted cyclopropene compounds with up to four different R groups which may be further substituted and Sisler teaches extensive substitution on the cyclopropene core.

Daly, in fact, teaches only a limited substitution pattern on the cyclopropene ring. Substituents are from one to four of hydrogen, C₁-C₄ alky, hydroxy, halogen, C₁-C₄ alkoxy, amino, and carboxy. No substituent group with greater than four atoms is disclosed or taught and the identity of the allowed groups is severely limited. Only one compound is exemplified, 1-methylcyclopropene. Daly teaches that the most preferable number of substituents is one (col. 9, lines 35-36) and the most preferable substituents are small groups (hydrogen, halogen, amino, carboxy), especially methyl.